



adelgids on bark

Univ. of Georgia



crawler



adelgids on bark

Univ. of Georgia



crawler

balsam woolly adelgid

Adelges piceae (Ratzeburg)

Adult adelgids are blackish purple, roughly spherical in shape, less than 1/32 inch (1 mm) long, and almost invisible to the naked eye. The adelgid produces a covering of white wax threads and appears as white, woolly dots about the size of pin heads on the surface of the tree's bole, limbs, and buds. Eggs are produced under the adults and are orange in color. The immature stage of the adelgid, known as a "crawler," is also orange, with legs and black eyes. Eggs and crawlers can be identified with the aid of a hand lens.

During the adelgid's feeding process, the host tree is stimulated to produce abnormal wood cells. This reduces the tree's ability to translocate food and water. Initial symptoms of adelgid attack may include gouting of buds or twig nodes and some twig and branch dieback. This is very evident on seedlings, young understory trees, and Christmas tree plantings. Other damage may be stunted shoot and needle growth and loss of apical dominance in natural stands. A heavily infested tree may die within 2 to 7 years. As the tree, dies, portions of the crown or the entire crown will turn red.

The adelgid has two generations per year, and occasionally three in the southern Appalachians. Eggs of the first generation hatch in late June and July, followed by a second generation in September and October. The "crawler" is the only mobile stage in the adelgid's life cycle. When a crawler begins feeding, it transforms into a first instar nymph and becomes stationary. Reproduction is parthenogenic; each female lays up to 200 eggs. The adelgid overwinters as a first instar nymph and continues its development in the spring when the host tree starts its annual growth cycle.

balsam woolly adelgid

Adelges piceae (Ratzeburg)

Adult adelgids are blackish purple, roughly spherical in shape, less than 1/32 inch (1 mm) long, and almost invisible to the naked eye. The adelgid produces a covering of white wax threads and appears as white, woolly dots about the size of pin heads on the surface of the tree's bole, limbs, and buds. Eggs are produced under the adults and are orange in color. The immature stage of the adelgid, known as a "crawler," is also orange, with legs and black eyes. Eggs and crawlers can be identified with the aid of a hand lens.

During the adelgid's feeding process, the host tree is stimulated to produce abnormal wood cells. This reduces the tree's ability to translocate food and water. Initial symptoms of adelgid attack may include gouting of buds or twig nodes and some twig and branch dieback. This is very evident on seedlings, young understory trees, and Christmas tree plantings. Other damage may be stunted shoot and needle growth and loss of apical dominance in natural stands. A heavily infested tree may die within 2 to 7 years. As the tree, dies, portions of the crown or the entire crown will turn red.

The adelgid has two generations per year, and occasionally three in the southern Appalachians. Eggs of the first generation hatch in late June and July, followed by a second generation in September and October. The "crawler" is the only mobile stage in the adelgid's life cycle. When a crawler begins feeding, it transforms into a first instar nymph and becomes stationary. Reproduction is parthenogenic; each female lays up to 200 eggs. The adelgid overwinters as a first instar nymph and continues its development in the spring when the host tree starts its annual growth cycle.